

RESULTS: DRIED BLOOD / URINE TEST

Accession #: 100035490 • Patient: JANE SMITH

Patient: JANE SMITH

Sex: Female

Age: 37 yr

Date of Birth: 1988-08-04

Accession #: 100035490



Sample received: 2026-09-05

Report issued: 2026-09-15

Health Care Professional: Jane Smith

Sample collection: 2026-03-01

HISTAMINE : DIAMINE OXIDASE (DAO)



* The reference range represents a quintile distribution, each quintile being 20 % of a population determined from archival data.

† Histamine values are determined from a Dried Urine sample. The red line at extreme left of bar represents Histamine values < 2.6 ng/mg Creat., and corresponds to values coincident with 'histapenia' in blood measurements.

Analysis performed by GB Diagnostics, 3704 General Patch Street NE, Albuquerque NM, USA - CLIA # 03D2182284.
The performance characteristics of this test were determined by GB Diagnostics. It has not been cleared by the U.S. Food and Drug Administration.

RESULTS: DRIED BLOOD / URINE TEST

Accession #: 100035490 • Patient: JANE SMITH

GENERAL COMMENTARY Histamine : Diamine Oxidase (DAO)

The comments provided here are for educational purposes only. The results in this report should not be interpreted as diagnostic, nor should they be viewed as treatment recommendations. Those decisions are the responsibility of the health care professional. Moreover, the reference ranges shown in this report are derived from a normal distribution of results, that encompass 95% of randomly selected individuals in a population (see below).

Histamine and Diamine Oxidase

Histamine is a nitrogenous compound known to be involved in 23 different physiological functions which include local immune responses, neurotransmission, as well as regulation of physiological function in the gastrointestinal tract.

Histamine is either stored, or rapidly metabolized and inactivated, by either of two degradative enzymes, histamine-N-methyltransferase or diamine oxidase (DAO); the latter being the principal enzyme observed in the digestive tract.

The testing of histamine, together with DAO levels, provides important information that is not revealed with standard food sensitivity tests. The latter is often thought to be the underlying cause of gut inflammation and dysfunction, when in fact the reason may actually be an imbalance of accumulated histamine and the capacity for its degradation by enzymes such as DAO. This imbalance may result in a condition known as Histamine Intolerance (HIT).¹¹ HIT is an acquired problem and is seen in approximately 3% of the population. Symptoms may occur in up to 20% of the population when histamine-rich food is consumed together with DAO inhibitors, such as alcohol. Women represent about 80% of those affected, with most of them aged 40 or over. It is important to note that the risk of developing this intolerance is increased in those individuals who suffer from inflammatory intestinal diseases or cross-sensitivities.¹¹

Why Test for Histamine?

Histamine is involved in many inflammatory and allergic processes, including both immediate and delayed hypersensitivity reactions. Histamine excess can be triggered by its release in the body as a result of a variety of environmental triggers, from the ingestion of foods with high histamine content, a deficiency in DAO, or both.

The Histamine reference range is a quintile distribution, extending from values of 2.6 ng/mg Creat to 250 ng/mg Creat. As histamine values approach the upper limit of the quintile range, there is an increasing probability of HIT.¹⁰ Urinary histamine levels can be influenced by recent dietary intake (histamine-rich foods) and potential urinary tract infections.

High levels of histamine (or elevated Histamine : DAO ratio)

- Runny nose, sneezing, congestion
- Itching, hives, skin flushing
- Dizziness or vertigo
- Headache, migraine
- Nausea, vomiting
- Intestinal cramps, gas
- Diarrhea

RESULTS: DRIED BLOOD / URINE TEST

Accession #: 100035490 • Patient: JANE SMITH

- Abnormal menstrual cycle
- Shortness of breath
- Abnormal heart rate
- High blood pressure
- Severe allergic reactions (anaphylaxis)
- Abnormalities may also arise in the following:
 - Memory
 - Body temperature
 - Circadian rhythm
 - Locomotion
 - Learning

What causes high histamine levels?

- Allergies (IgE reactions)
- Gluten Intolerance
- Small Intestinal Bacterial Overgrowth (SIBO)
- Intestinal Permeability ("Leaky Gut")
- Gastrointestinal bleeding
- Histamine-rich foods
- DAO deficiency or DAO-blocking foods: alcohol, energy drinks, and tea
- Genetic mutations (common in people of Asian descent)
- Inflammatory bowel diseases: Crohn's, ulcerative colitis
- Medications:
 - Non-steroidal anti-inflammatory drugs (eg: Ibuprofen-Motrin, ASA-Aspirin)
 - Antidepressants (eg: Effexor, Zoloft, Prozac, Cymbalta)
 - Immune modulators (eg: Enbrel, Humira)
 - Anti-arrhythmics (eg: Propranolol, Norvasc, Cardizem)
 - H2 blockers (eg: Zantac, Pepcid, Tagamet)

Very low levels of histamine:

- Fatigue
- Sleep-wake disorders
- Depression and anxiety in older adults; paranoia in younger people
- Convulsions

What causes low histamine levels?

A low histamine diet can cause a significant, sometimes 8-fold, reduction in urinary histamine levels. Reduced urinary excretion of histamine, and its metabolites, can occur due to renal impairment. Also, the overuse of H1 receptor blockers (antihistamines) can sometimes result in very low levels of histamine.

Excess copper can create low levels by decreasing histamine in the brain. In turn, the lowered levels of histamine allow more copper to accumulate. High copper in the brain may lead to a state of restlessness, insomnia, violence, depression, irritability, paranoia, and high blood pressure.

RESULTS: DRIED BLOOD / URINE TESTAccession #: 100035490 • Patient: JANE SMITH

Why Test for DAO?

The ingestion of histamine rich food, alcohol or drugs that release histamine or block DAO, may provoke an imbalance of accumulated histamine and the capacity for its degradation, already referred to above as Histamine Intolerance (HIT).

An impaired histamine degradation, based on reduced DAO activity and the resulting histamine excess, may cause numerous symptoms mimicking an allergic reaction.¹¹

DAO activity does not depend on the DAO alone, but also on cofactors such as vitamin C, vitamin B6, copper or manganese ions. Copper is a central component of DAO. A deficiency in copper can result in insufficient DAO being produced. Vitamin B6 is a cofactor of DAO. If vitamin B6 is missing, DAO is unable to degrade histamine.

Therefore, in assessing HIT via the DAO activity test, one should also consider determining the levels of these cofactors. The symptoms of HIT can be caused by low DAO activity because the above-mentioned cofactors are not sufficiently available.

If the DAO levels are in the normal range but the histamine levels are high, it may indicate that the issue is not insufficient DAO, but rather an overproduction of histamine, due to factors such as gut dysbiosis.

However, if the histamine levels are normal, but the DAO levels are very low, it suggests a possible genetic deficiency of diamine oxidase.

DAO results, in this report, are obtained from a Dried Blood Spot sample.

The DAO reference range is a quintile distribution, extending from low values of < 30 ng/g Hgb to 4000 ng/g Hgb. The DAO quintile reference range has no significant daily variations or gender differences. As levels go from 4000 to < 100 ng/g Hgb, it indicates an increasing probability of HIT.¹¹

Low levels of DAO:

- Skin rash and pruritis (itching), urticaria (hives), eczema, psoriasis
- Nasal congestion, asthma
- Headache, migraine
- Chronic fatigue
- Anxiety, depression
- Inflammation, irritable bowel syndrome (IBS)
- Estrogen dominance, dysmenorrhea, Premenstrual Syndrome (PMS)
- Muscular pain, fibromyalgia
- Rheumatoid arthritis
- Hypertension, hypotension, arrhythmia,
- Multiple sclerosis and other neurological conditions

RESULTS: DRIED BLOOD / URINE TEST

Accession #: 100035490 • Patient: JANE SMITH

Determination of DAO activity, together with a detailed history, helps to differentiate food allergy and histamine intolerance. It should be performed in suspected patients who have headaches, urticaria, pruritus, diarrhea and hypotension, where food allergy has been excluded.¹²

Individuals who are unable to metabolize histamine will often improve with a variety of antihistamines. Because DAO formation occurs in the gastrointestinal system, lower than normal levels are suggestive of poor digestive dysfunction, as well as problems in the intestinal barrier.¹³

The Importance of the Histamine : DAO Ratio

The Histamine : DAO Ratio is helpful in highlighting the imbalances in histamine and DAO levels.

High Ratio: There is a relatively high level of free histamine in the system from both endogenous and exogenous sources, and/or there is an insufficient amount of DAO enzyme available for histamine degradation. It should be noted that even if DAO levels are normal, symptoms may still occur if the histamine levels are very high. As levels reach into the elevated range, histamine intolerance should be considered.

Low Ratio: The amount of free histamine is relatively low, and/or there is sufficient DAO enzyme available to degrade the histamine that is present. An adequate intake of healthy fats and other nutrients, like phosphorus, zinc, magnesium, iron, and vitamin B12, are known to play a role in enhancing DAO activity.¹⁴ Eating mainly low-histamine foods may reduce exposure to histamine and lower its accumulation in the body. This combination of factors will lead to a decrease in the Histamine : DAO ratio.

DAO: A Biomarker of Intestinal Barrier Integrity

The degradative enzyme, Diamine Oxidase (DAO), is the principal enzyme observed in the digestive tract which scavenges extracellular histamine. Recent research has begun to shed light on another important aspect of DAO activity, which is unique among intestinal mucosal enzymes: Circulating blood levels of DAO represent a reliable marker of mucosal maturation & integrity¹⁵. Indeed, serum DAO activity has been shown to correlate with intestinal permeability of the small intestine.¹⁶

Intestinal Barrier Changes

Intestinal barrier damage is due to a number of potential causes, amongst them mechanical, microbial and/or enzymatic action. These vectors of damage can be due to:

- Genetic predisposition
- Dysbiosis
- Infections
- Loss of luminal mucous barrier, with a concomitant decrease in levels of SIgA and antimicrobial peptides (AMPs).

RESULTS: DRIED BLOOD / URINE TEST

Accession #: 100035490 • Patient: JANE SMITH

• High luminal histamine levels from endogenous or exogenous sources, or both. Increased histamine can occur due to gut microbial action, a diet rich in histamine or histidine, an increased mast cell response as part of an immune reaction and acute or chronic stress.

DAO is synthesized by mucosal cells of intestinal villi crypts. The damage caused by one or more of the factors above can lead to a loss of barrier integrity, which in turn can lead to further inflammatory reactions. All of this will often lead to increased intestinal permeability, which will then enter into a vicious cycle of more inflammation and barrier damage. During the period of initial damage, the mucosal cells release increased amounts of DAO which increases its serum concentration. Therefore, a change in blood DAO concentration is an indication of damage to the intestinal cavity. High DAO in blood is tightly linked to abnormal intestinal barrier function in an acute stage.¹⁷

With more chronic damage, as often seen in cases of inflammatory bowel diseases (IBD), such as Chron's disease or Ulcerative Colitis, one begins to encounter apoptosis of the crypt cells within the villi. This decrease or loss of DAO levels in the gut lumen & blood is a sign of chronic or late stage mucosal damage to the barrier, and is a sign of breakdown in cell architecture. More specifically, it is a decrease in DAO production capacity due to the destroyed or dysfunctional specialized mucosal cells¹⁶. This makes DAO a sensitive & accurate marker for monitoring Crohn's Disease activity and other inflammatory bowel conditions.

The Histamine : DAO Ratio and Intestinal Permeability

As stated previously, the Histamine : DAO Ratio is helpful in highlighting the imbalances in histamine and DAO levels as well as highlighting changes in intestinal barrier integrity and permeability. Abnormal ratios have been found to highly correlate with many common gut-related issues and other complaints.

High Ratio: A high ratio is an indication that there is a relatively high level of free histamine in the system, often combined with insufficient DAO enzyme available for histamine degradation. When histamine levels are 'normal', and yet the ratio to DAO is still high, it is often a sign of abnormally low production of DAO, due to a breakdown in cell architecture, or even cell destruction (crypt hyperplasia), as seen in cases of IBD and increased intestinal permeability. DAO co-factors may also need to be considered

Low Ratio: When high levels of the DAO enzyme are available to degrade normal or low amounts of free Histamine, it could be a sign of early breakdown in barrier integrity, as DAO is often elevated in the initial stages of mucosal barrier dysfunction.

References

1. Vanuytsel T et al. Tissue Barriers 2013; 1: 1-9; 2. Fasano A et al. Lancet 2000; 355: 1518-19; 3. Fasano A. Ann N Y Acad Sci 2012; 1258: 25-33; 4. Sapone A et al. Diabetes 2006; 55: 1443-1449; 5. Pacifico, L et al. World J of Gastroenterol 2014; 20: 17107-17114; 6. Skardelly M, et al. Transl Oncol 2009; 2: 117-120; 7. Rittirsch D, et al. Am J Physiol Lung Cell Mol Physiol 2013; 304: L863-L872; 8. Turner JR. Am J Pathol 2006; 169: 1901-1909; 9. Lee SH. Intest Res 2015; 13: 11-18; 10. Jean Dyer BS et al. J Allergy Clin Immunol 1982; 70 (2): 82-87.; 11. Maintz L, Novak N. Am J Clin Nutr 2007; 85:1185-1196; 12. Music E, et al. Food Allergy and Anaphylaxis Meeting 2011, Venice, Italy. 17-19 February, 2011; 13. Farhadi A, et al. J Gastroenterol and Hepatol 2003; 18: 479-497.; 14. Miyoshi et al. Nutrition. 2017; 39-40: 67-70. 15. Luk et al. J Clin Invest 1980; 66: 66-70.; 16. Honzawa et al. Inflamm Bowel Dis 2011; 17: Letter to Editor.; 17. Cai et al. Gastroenterology Research and Practice Vol 2019; 18. . Murch O et al. Intensive Care Med 2007; 33:13-24.; 19. Mousa H et al. Biochemistry and Biophysics Reports 2023; 34: 101445.; 20. Tobias PS, Ulevich RJ. Immunobiology 1993; 187: 227-232.; A. Lian P et al. Nutrients 2020; 12: 734; B. Sun M et al. Cell Death Discovery 2024; 10: 455; C. Ural D A. IJVAR 2022; 5(2): 47-49.



[Back to Control Panel >>](#)

RESULTS: DRIED BLOOD / URINE TEST

Accession #: 100035490 • Patient: JANE SMITH
